

**Committed For A Better Life** 

# CRITICAL CARE

12 1

Dedicated to Provide Healthy Life With Excellent Quality.

# COMPANY Profile 2022

MEDISUM BIOTECH PVT. LTD. was founded in march 2022 with the aim of making affordable healthcare reachable to every Indian household.

Medisum biotech having its registered corporate office in Delhi is a balance of experience and zeal with its chairman having an experience of 40 years in the pharma industry and young directors thrieving to make thier mark.

Our focus here at medisum is at providing excellant quality medicines at an affordable price to every citizen in the country.

The key forces driving our company are innovation, integrity & service to the human kind.

# Committed for a better life.



**Certificate of Registration** 

This Certificate certify that the Management System of

# **MEDISUM BIOTECH PRIVATE LIMITED**

E-5/16, Krishna Nagar, Delhi - 110051, India.

has been audited by ICV and found to be in compliance with the requirements of the standard

ISO 9001:2015

(Quality Management System)

# This certificate is valid for the following scope:

Manufacturer, Trader, Dealers, Job Workers, Processors, Sellers, Retailer, Buyers, Wholesalers, Importers, Exporters in Pharmaceuticals, Pharmaceutical-fine-Chemicals, Pharmaceutical Products and Bulkdrugs etc.

Certificate No: IN/18907689/4540Initial Registration Date: 28/03/2023Certificate Issue Date: 28/03/2023Certification Expiry/Re Certification Date: 27/03/2026







CEO ICV ASSESSMENTS PVT. LTD.

25, First Floor, Prabhu Complex, Shiva Market, Pitampura New Delhi - 110034 +91 - 11-41548315, info@icvassessments.com, www.icvassessments.com

Validity of this Certificate is subject to completion of surveillance audit on or before of due date. In case Surveillance audit not allowed to be conducted this Certificate shall be suspended / withdrawal.
This Certificate is intellectual Property of ICV and Certificate should be returned to ICV in case of non compliance of certification procedure. Authenticity of this certificate can be verified at www.icvassessments.com

# **CERTIFICATE** OF REGISTRATION



This is to certify that the management system of

# **MEDISUM BIOTECH PRIVATE LIMITED**

has been formally assessed by INTERNATIONAL CERTIFICATION & INSPECTION UK LTD. and found to comply with the requirements of

# id found to comply with the requirements

# GMP

# (Good Manufacturing Practice)

# Scope of Registration:

Manufacturer, Trader, Dealers, Job Workers, Processors, Sellers, Retailer, Buyers, Wholesalers, Importers, Exporters in Pharmaceuticals, Pharmaceutical-fine-Chemicals, Pharmaceutical Products and Bulkdrugs etc.

# **Registered Site (s):**

E-5/16, Krishna Nagar, Delhi - 110051, India.

# :: Certificate No :: ICI/3128543/23

Date of initial registration:

21 April 2023

First Surveillance Audit on or before:

21 March 2024

Second Surveillance Audit on or before:

21 March 2025

**Re-certification Due:** 

20 April 2026

This Certificate is property of ICI UK Ltd. and remains valid subject to satisfactory surveillance audits.

Peter Collin Ex. Director

International Certification & Inspection UK Limited. Kemp House, 160 City Road, London, EC1V 2NX, United Kingdom





To check validity of the certificate please visit at www.ici-uk.co

This certification of registration is issued by International Certification & Inspection UK Limited accredited with Accreditation Board For Certification bodies (www.ab-cb.org). This certificate remains the property of International Certification & Inspection UK Limited and must be returned upon request.



# Alendrone-70 TABLETS

(Each Uncoated Tablet Contains: Sodium Alendronate Trihydrate USP Eq.to Alendronic Acid 70mg + Excipients q.s.)



#### Indication:

► Alendronate is used to prevent and treat certain types of bone loss (osteoporosis) in adults. Osteoporosis causes bones to become thinner and break more easily.

#### Dosage:

Directed by the physician

# Alfuze-D TABLETS

Each Film-coated tablet contains : Alfuzosin Hydrochloride IP 10mg. + Dutasteride IP 0.5mg. Expients q.s.Colour : Titanium Dioxide IP



#### Indication:

 Alfuzosin +Dutasteride is a combination of two medicines: Alfuzosin and Dutasteride, which relieves the symptoms of benign prostatic hyperplasia. Alfuzosin is an alpha-blocker. It works by relaxing the muscle around the bladder exit and prostate gland so as to allow easy passage of urine

### Dosage:

Directed by the physician

# Alfuze-10 TABLETS

Each Film-coated prolonged release tablet containsn Alfuzosin Hydrochloride IP 10mg. Excipients q.s. colour : Titanium dioxide IP





#### Indication:

 Alfuzosin is used to treat the signs and symptoms of benign enlargement of the prostate (benign prostatic hyperplasia or BPH). Benign enlargement of the prostate is a problem that can occur in men as they get older. The prostate gland is located below the bladder.

#### Dosage:

Directed by the physician

# Apexbn-2.5 TABLETS

Each Film Coated Tablet Contains: Apixaban 2.5 Mg



#### Indication:

- Apixaban is indicated for reducing the risk of stroke
- Apixaban is a type of medicine known as an anticoagulant.



# Apexbn-5 TABLETS

Each Film Coated Tablet Contains: Apixaban 5 Mg



#### Indication:

- Apixaban is indicated for reducing the risk of stroke
- Apixaban is a type of medicine known as an anticoagulant.

#### Dosage:

Directed by the physician

# Atova–Sure 20 TABLETS

Each film coated tablet contains: Atorvastatin Calcium IP Eq. to Atorvastatin 20 mg + Excipients q.s.



#### Indication:

- Reduce the risk of heart attack and stroke.
- Helpful as on aggressive lipid lowering therapy.

#### Dosage:

Directed by the physician

# Atova–Sure 10 TABLETS

Each film coated tablet contains: Atorvastatin Calcium IP Eq. to Atorvastatin 10 mg + Excipients q.s.



#### Indication:

- Reduce the risk of heart attack and stroke.
- Helpful as on aggressive lipid lowering therapy.

#### Dosage:

Directed by the physician

# Atova–Sure 40 TABLETS

Each film coated tablet contains: Atorvastatin Calcium IP Eq. to Atorvastatin 40 mg + Excipients q.s.



#### Indication:

- Reduce the risk of heart attack and stroke.
- Helpful as on aggressive lipid lowering therapy.



# Azathiomed-50 TABLETS

Each Uncoated Tablet Contains: Azathioprine IP 50 mg



#### Indication:

Azathioprine is used to prevent organ rejection in people who have received a kidney transplant. It is usually taken along with other medications to allow your new kidney to function normally. Azathioprine is also used to treat rheumatoid arthritis.

#### Dosage:

Directed by the physician

### Bisolol-5 TABLETS

Each Uncoated tablet contains: Bisoprolol Fumarate USP 2.5 mg + Excipients q.s.



#### Indication:

- Bisoprolol is used to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.
- ► Bisoprolol is used to lowers the heart rate, blood pressure, and strain on the heart. **Dosage:**

#### Directed by the physician

# Bisolol-2.5 TABLETS

Each Uncoated tablet contains: Bisoprolol Fumarate USP 2.5 mg + Excipients q.s.



#### Indication:

- Bisoprolol is used to treat high blood pressure (hypertension).
   Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.
- ► Bisoprolol is used to lowers the heart rate, blood pressure, and strain on the heart.

#### Dosage:

Directed by the physician

### **Baclofine-10** TABLETS

Each Uncoated tablet contains: Baclofen IP 10 mg + Excipients q.s.



#### Indication:

 Baclofen is used to help relax certain muscles in your body. It relieves the spasms, cramping, and tightness of muscles caused by medical problems, including multiple sclerosis or certain injuries to the spine



# Carvelol-3.125 TABLETS

Each Uncoated Tablet Contains: Carvedilol IP 3.125 mg + Excipients q.s.



#### Indication:

This drug works by blocking the action of certain natural substances in your body, such as epinephrine, on the heart and blood vessels. This effect lowers your heart rate, blood pressure, and strain on your heart.

#### Dosage:

Directed by the physician

# Carvelol-12.5 TABLETS

Each Uncoated Tablet Contains: Carvedilol IP 12.5 mg + Excipients q.s.



#### Indication:

This drug works by blocking the action of certain natural substances in your body, such as epinephrine, on the heart and blood vessels. This effect lowers your heart rate, blood pressure, and strain on your heart.

#### Dosage:

Directed by the physician

# Carvelol-6.25 TABLETS

Each Uncoated Tablet Contains: Carvedilol IP 6.25 mg + Excipients q.s.



#### Indication:

This drug works by blocking the action of certain natural substances in your body, such as epinephrine, on the heart and blood vessels. This effect lowers your heart rate, blood pressure, and strain on your heart.

#### Dosage:

Directed by the physician

# Carvelol-CR 10 TABLETS

Each Uncoated Tablet Contains: Carvedilol IP 12.5 mg + Excipients q.s.



#### Indication:

This drug works by blocking the action of certain natural substances in your body, such as epinephrine, on the heart and blood vessels. This effect lowers your heart rate, blood pressure, and strain on your heart.

#### Dosage:



# Clindimed-5 TABLETS

Each film coated tablet contains: Cilnidipine Tablets IP 5mg



#### Indication:

Cilnidipine is used in the treatment of Hypertension (high blood pressure), Angina (heart-related chest pain), heart attack and stroke. Cilnidipine is a calcium channel blocker. It lowers blood pressure by relaxing blood vessels, which makes the heart more efficient at pumping blood throughout the body. Dosage:

Directed by the physician

# Clopisure-75 TABLETS

Each Film Coated Tablet contains: Clopidogrel Bisulphate IP eq. to Clopidogrel 75 mg + Excipients q.s.



#### Directed by the physician

# Clindimed-10 TABLETS

Each film coated tablet contains: Cilnidipine Tablets IP 10mg



#### Indication:

Cilnidipine is used in the treatment of Hypertension (high blood pressure), Angina (heart-related chest pain), heart attack and stroke. Cilnidipine is a calcium channel blocker. It lowers blood pressure by relaxing blood vessels, which makes the heart more efficient at pumping blood throughout the body.

#### Dosage:

Directed by the physician

# **Clopisure-AP** TABLETS

Each Film Coated Tablet contains: Clopidogrel Bisulphate IP eq. to Clopidogrel 75 mg +Aspirin 75mg Excipients q.s.



#### Indication:

- Enhance antithrombotic
- Reduce the risk of stroke & MI
- Inhibits platelet aggregation

# Dosage:



# Chloridone-6.25 TABLETS

Each unoated Tablet contains: Chlorthalidone IP 6.25 Excipients q.s.



#### Indication:

Chlorthalidone which belongs to the group of medicines called Diuretics. It is used to treat hypertension (high blood pressure) and heart failure. It also reduces fluid accumulation (edema) from the body that is caused due to certain liver or kidney diseases. Dosage:

Directed by the physician

# Colchisure-0.5 TABLETS

Each Uncoated Tablet Contains: Colchicine IP 0.5 mg + Excipients q.s.



Directed by the physician

# Chloridone-12.5 TABLETS

Each unoated Tablet contains: Chlorthalidone IP 12.5 Excipients q.s.



#### Indication:

► Chlorthalidone which belongs to the group of medicines called Diuretics. It is used to treat hypertension (high blood pressure) and heart failure. It also reduces fluid accumulation (edema) from the body that is caused due to certain liver or kidney diseases.

#### Dosage:

Directed by the physician

### **Dapasum-5** TABLETS

Each Film Coated tablet Contains: Dapagliflozin propanediol Monohydrate Eq.to Dapagliflozin 5mg + Excipients q.s



#### Indication:

 Used for treatment of type 2 diabeties. Helpful in treatment of chronic kidney disease with Sugar.

# Dosage:



# Dapasum-10 TABLETS

Each Film Coated tablet Contains: Dapagliflozin propanediol Monohydrate Eq.to Dapagliflozin 10mg + Excipients q.s



 Used for treatment of type 2 diabeties. Helpful in treatment of chronic kidney disease with Sugar.

#### Dosage:

Directed by the physician

# Dapasum-M 1000 TABLETS

Each Film Coated Tablet Contains: Dapagliflozin (10mg) + Metformin (1000mg) + Excipients q.s



#### Indication:

 Dapagliflozin works in the kidneys to prevent absorption of glucose (blood sugar). This helps lower the blood sugar level. Metformin reduces the absorption of sugar from the stomach, reduces the release of stored sugar from the liver, and helps your body use sugar better.

#### **Dosage:** Directed by the physician

# Dapasum-M 500 TABLETS

Each Film Coated Tablet Contains: Dapagliflozin (10mg) + Metformin (500mg) + Excipients q.s



#### Indication:

 Dapagliflozin works in the kidneys to prevent absorption of glucose (blood sugar). This helps lower the blood sugar level. Metformin reduces the absorption of sugar from the stomach, reduces the release of stored sugar from the liver, and helps your body use sugar better.


#### Dosage:

Directed by the physician

### Donzy-5 TABLETS

Each Film Coated Tablet Contains: Donepezil Hydrochloride IP 5 mg + Excipients q.s





### Indication:

 DONEPEZIL treats memory loss and confusion (dementia) in people who have Alzheimer disease. It works by improving attention, memory, and the ability to engage in daily activities.

#### Dosage:



# Donzy-10 TABLETS

Each Film Coated Tablet Contains: Donepezil Hydrochloride IP 10 mg + Excipients q.s



#### Indication:

 DONEPEZIL treats memory loss and confusion (dementia) in people who have Alzheimer disease. It works by improving attention, memory, and the ability to engage in daily activities.

#### Dosage:

Directed by the physician

### Enasum--5 TABLETS

Each Film Coated tablet Contains: Each Enalapril Maleate Tablets IP 5mg



Enalapril is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

#### Dosage:

Directed by the physician

# **Dydromed-10** TABLETS

(each Film Coated Tab Contains Dydrogesterone I.P. 10 Mg + Excipients Q.s.)

# 10x1x10



#### Indication:

Dydrogesterone helps to regulate the healthy growth and normal shedding of the uterus lining. Therefore, it may be useful in the treatment of menstrual disorders such as absent, irregular or painful menstrual periods, infertility, premenstrual syndrome and endometriosis.

#### Dosage:

Directed by the physician

### Enasum-2.5 TABLETS

Each Film Coated tablet Contains: Each Enalapril Maleate Tablets IP 2.5mg



#### Indication:

Enalapril is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

#### Dosage:



# Enaveer-0.5 TABLETS

Each Film Coated Tablet Contains: Entecavir Tablets 0.5mg. Excipients g.s.



#### Indication:

► Entecavir is used to treat liver infection caused by hepatitis virus. It belongs to the family of medicines called antivirals. Antivirals are used to treat infections that are caused by viruses. This medicine will not cure the hepatitis B virus, but it will keep it from reproducing and causing more liver damage. Dosage:

Directed by the physician

# Etocoxum-90 TABLETS

Each Film coated tablet contains Etoricoxib IP 90 mg



► Etoricoxib helps to reduce the pain and swelling (inflammation) in the joints and muscles of people 16 years of age and older with osteoarthritis, rheumatoid arthritis, ankylosing spondylitis and gout.

#### Dosage:

Directed by the physician

# Ezimib-10 TABLETS

Each uncoated tablet contains : Ezetimibe IP 10mg. Excipients q.s.



Ezetimibe is a dyslipidemic agent used to treat people with hyperlipidemia. It was FDA-approved in 2002. Ezetimibe is the most commonly used nonstatin agent, which lowers LDL-C levels by 13% to 20%.[1] Ezetimibe is an inhibitor of intestinal cholesterol absorption.

#### Dosage:

Directed by the physician

# Etocoxum-120 TABLETS

Each Film coated tablet contains Etoricoxib IP 120 mg



• Etoricoxib helps to reduce the pain and swelling (inflammation) in the joints and muscles of people 16 years of age and older with osteoarthritis, rheumatoid arthritis, ankylosing spondylitis and gout.

# Dosage:



# Feronum-200 TABLETS

Each Film Coated Tablet Contains: Faropenem Sodium 200 Hydrate eq. to Faropenem 200 mg + Excipients q.s



#### Indication:

- ▶ Well tolerated and has a favourable safety profile
- ► Exhibits favourable effect on nasal congestion

#### Dosage:

Directed by the physician

# Flucozole TABLETS

Each uncoated tablet contains: Fluconazole Tablet IP 150mg. Excipients q.s.



#### Indication:

Fluconazole is used to treat serious fungal or yeast infections, including vaginal candidiasis, oropharyngeal candidiasis (thrush, oral thrush), esophageal candidiasis (candida esophagitis), other candida infections (including urinary tract infections, peritonitis [inflammation of the lining of the stomach], and infections that may occur in different parts of the body), or fungal (cryptococcal) meningitis.

Dosage:

Directed by the physician

# Feronum-300 SR TABLETS

Each Film Coated Sustained Release Tablet Contains: Faropenem Sodium 300 Hydrate eq. to Faropenem 300 mg + Excipients q.s

10x1x6



#### Indication:

- ▶ Well tolerated and has a favourable safety profile
- Exhibits favourable effect on nasal congestion

#### Dosage:

Directed by the physician

# Gabasure-NT (100/10) TABLETS

Each film coated tablet contains: Gabapentin IP 100mg + Nortriptyline Hydrochloride IP 10 mg, Eq. to Nortriptyline, Excipients q.s.





# **Glicosum-1** TABLETS

Each uncoated tablet contains: Glimepiride IP 1 mg + Excipients q.s.



- Very effective as initial drug theray
- ▶ Less prone to cause hypoglycemia

#### Dosage:

Directed by the physician

# Glicosum-M1 SR TABLETS

Each uncoated bilayered tablet contains: Glimepiride IP 1 mg + Metformin Hydrochloride IP 500 mg (In Sustained Release Form) + Excipients q.s.



#### Indication:

- Very effective as initial drug theray
- Less prone to cause hypoglycemia

Dosage: Directed by the physician

### **Glicosum-2** TABLETS

Each uncoated tablet contains: Glimepiride IP 2 mg + Excipients q.s.





#### Indication:

- Very effective as initial drug theray
- ▶ Less prone to cause hypoglycemia

#### Dosage:

Directed by the physician

# Glicosum-M2 SR TABLETS

Each uncoated bilayered tablet contains: Glimepiride IP 2 mg + Metformin Hydrochloride IP 500 mg (In Sustained Release Form) + Excipients q.s.



#### Indication:

- Very effective as initial drug theray
- Less prone to cause hypoglycemia





# Glicosum-M1 Forte TABLETS

Each uncoated bilayered tablet contains: Glimepiride IP 1 mg + Metformin Hydrochloride IP 1000 mg (In Sustained Release Form) + Excipients q.s.



#### Indication:

- Very effective as initial drug theray
- ▶ Less prone to cause hypoglycemia

#### Dosage:

Directed by the physician

# Glicosum-PG2 TABLETS

Each uncoated bilayered tablet contains : Glimepiride IP 2 mg + Pioglitazone Hydrochloride IP Eq. to Pioglitazone 15 mg + Metformin Hydrochloride IP 500 mg (In Extended Release Form) + Excipients q.s.



- Less prone to cause hypoglycemia

### Dosage:

Directed by the physician

# **Glicosum-M2 Forte** TABLETS

Each uncoated bilayered tablet contains: Glimepiride IP 2 mg + Metformin Hydrochloride IP 1000 mg (In Sustained Release Form) + Excipients q.s.



#### Indication:

- Very effective as initial drug theray
- ▶ Less prone to cause hypoglycemia

#### Dosage:

Directed by the physician

# Gabasure-NT (400/10) TABLETS

Each film coated tablet contains: Gabapentin IP 400mg + Nortriptyline Hydrochloride IP 10 mg, Eq. to Nortriptyline, Excipients q.s.



- Post therapy Neuraliga
- Drug induced neuropathy





# Glizomed-40

TABLETS

Each Uncoated Tablet Contains: Gliclazide IP 40 mg + Excipients q.s.



#### Indication:

Gliclazide is used with a proper diet and exercise program to control high blood sugar in people with type 2 diabetes. controlling high blood sugar helps prevent heart disease, strokes, kidney disease, blindness, circulation problems, and decreased sexual ability.

#### Dosage:

Directed by the physician

# Glizomed 30 MR

TABLETS

Each Uncoated Modified Release Tablet Containes: Gliclazide IP 30 mg + Excipients q.s.




#### Indication:

 Gliclazide is used with a proper diet and exercise program to control high blood sugar in people with type 2 diabetes. controlling high blood sugar helps prevent heart disease, strokes, kidney disease, blindness, circulation problems, and decreased sexual ability.

**Dosage:** Directed by the physician

# Glizomed-80 TABLETS

Each Uncoated Tablet Contains: Gliclazide IP 80 mg + Excipients q.s.



#### Indication:


Gliclazide is used with a proper diet and exercise program to control high blood sugar in people with type 2 diabetes. controlling high blood sugar helps prevent heart disease, strokes, kidney disease, blindness, circulation problems, and decreased sexual ability.

#### Dosage:

Directed by the physician

### Glizomed 60 MR TABLETS

Each Uncoated Modified Release Tablet Containes: Gliclazide IP 60 mg + Excipients q.s.



Gliclazide is used with a proper diet and exercise program to control high blood sugar in people with type 2 diabetes. controlling high blood sugar helps prevent heart disease, strokes, kidney disease, blindness, circulation problems, and decreased sexual ability.

#### Dosage:



#### Glizomed M 80 TABLETS

Each film tablet contains: Gliclazide IP80 mg + Metformin Hydrochloride IP500 mg + Excipients q.s.



#### Indication:

▶ Gliclazide and Metformin. This medicine is used in the treatment of type 2 diabetes mellitus (DM). It improves blood glucose levels in adults when taken along with proper diet and regular exercise.

#### Dosage:

Directed by the physician

# Ivadine-5 TABLETS

Each Film Coated Tablet Contains: Ivabradine Hydrochloride Eq. to Ivabradine 5 mg + Excipients q.s.



#### Indication:

- Works by affecting heart's electrical activity in order to slow the heart rate
- ▶ Used in adults with chronic heart failure

#### Dosage: Directed by the physician

# Indomide-1.5SR TABLETS

Each film coated sustained release tablet contains: Indapamide IP 1.5 + Excipients q.s



#### Indication:

▶ Indapamide, a 'water pill,' is used to reduce the swelling and fluid retention caused by heart disease. It also is used to treat high blood pressure. It causes the kidneys to get rid of unneeded water and salt from the body into the urine.

#### Dosage:

Directed by the physician

# Labetalod-100 TABLETS

Each Uncoated Tablet Contains: Labetalol Hydrochloride IP 100 mg



#### Indication:

► Labetalol maintiains cardiac output and also maintains peripheral, cerebral, renal and coronary blood flow



# Lacosamed-50 TABLETS

Each Film Coated Tablet Contains: Lacosamide 50 mg + Excipients q.s.



#### ndication:

▶ Lacosamide is used to treat partial-onset seizures. It is also used with other medicines to treat primary generalized tonic-clonic seizures. It acts on the central nervous system (CNS) to reduce the number and severity of seizures.

#### osage:

Directed by the physician

# Lin-Gip-5 TABLETS

Linagliptin 5mg.



#### Indication:

- Linagliptin is a medicine used to treat type 2 diabetes.
- ▶ Type 2 diabetes is a condition where the body does not
- make enough insulin, or the insulin that it makes does not work properly. This can cause high blood sugar levels (hyperglycaemia).

#### Dosage:

Directed by the physician

# Lacosamed-100 TABLETS

Each Film Coated Tablet Contains: Lacosamide 100 mg + Excipients q.s.



#### Indication:

▶ Lacosamide is used to treat partial-onset seizures. It is also used with other medicines to treat primary generalized tonic-clonic seizures. It acts on the central nervous system (CNS) to reduce the number and severity of seizures.

#### **Dosage:**

Directed by the physician

# Lin-Gip-M 500 TABLETS

Linagliptin 2.5mg., Metformin-500mg.



#### Indication:

- Linagliptin is a medicine used to treat type 2 diabetes.
- ▶ Type 2 diabetes is a condition where the body does not make enough insulin, or the insulin that it makes does not work properly.
- This can cause high blood sugar levels (hyperglycaemia).

#### Dosage:





# Lin-Gip-M 1000 TABLETS

Linagliptin 2.5mg., Metformin-1000mg.



#### Indication:

- ▶ Linagliptin is a medicine used to treat type 2 diabetes.
- ▶ Type 2 diabetes is a condition where the body does not
- make enough insulin, or the insulin that it makes does not work properly.This can cause high blood sugar levels (hyperglycaemia).
- Dosage:

Directed by the physician

# Livo Ctram-500 TABLETS

Each film coated tablet Contains : Levetiracetam IP 500 mg + Excipients q.s.



#### Indication:

 Levetiracetam is a novel antiepileptic drug used to treat partial, myoclonic, and tonic-clonic seizures. In 2000, the FDA approved the use of the oral formulation as adjunctive therapy for the treatment of focal seizures, myoclonic seizures, and primary generalized seizures.

#### Dosage:

Directed by the physician

# Livo Ctram-250 TABLETS

Each film coated tablet Contains : Levetiracetam IP 250 mg + Excipients q.s.



#### Indication:

Levetiracetam is a novel antiepileptic drug used to treat partial, myoclonic, and tonic-clonic seizures. In 2000, the FDA approved the use of the oral formulation as adjunctive therapy for the treatment of focal seizures, myoclonic seizures, and primary generalized seizures.

#### Dosage:

Directed by the physician

# Livo Ctram-750 TABLETS

Each film coated tablet Contains : Levetiracetam IP 750 mg + Excipients q.s.



#### Indication:

 Levetiracetam is a novel antiepileptic drug used to treat partial, myoclonic, and tonic-clonic seizures. In 2000, the FDA approved the use of the oral formulation as adjunctive therapy for the treatment of focal seizures, myoclonic seizures, and primary generalized seizures.

#### Dosage:



# Livo Ctram-100 TABLETS

Each film coated tablet Contains : Levetiracetam IP 1000mg + Excipients q.s.



#### Indication:

 Levetiracetam is a novel antiepileptic drug used to treat partial, myoclonic, and tonic-clonic seizures. In 2000, the FDA approved the use of the oral formulation as adjunctive therapy for the treatment of focal seizures, myoclonic seizures, and primary generalized seizures.

#### Dosage:

Directed by the physician

#### Losarsum-25

TABLETS

Each film coated tablet contains: Losartan Potassium IP 25 mg + Excipients q.s.



 Losartan is used to treat high blood pressure (hypertension) and to help protect the kidneys from damage due to diabetes. It is also used to lower the risk of strokes in patients with high blood pressure and an enlarged heart.

protect the kidneys from damage due to diabetes. It is also used to lower

the risk of strokes in patients with high blood pressure and an enlarged


#### Dosage:

heart.

Dosage:

Directed by the physician

Directed by the physician



Losartan is used to treat high blood pressure (hypertension) and to help protect the kidneys from damage due to diabetes. It is also used to lower the risk of strokes in patients with high blood pressure and an enlarged heart.

#### Dosage:



# Mesalamed-400 TABLETS

Each Delayed Release Tablet Contains: Mesalamine IP 400 mg + Excipients q.s.



#### Indication:

Mesalamine is used to treat and prevent flare-ups of mild to moderately active ulcerative colitis (an inflammatory bowel disease). It works inside the bowels to reduce inflammation and other symptoms of the disease. This medicine is available only with your doctor's prescription.

#### Dosage:

Directed by the physician

# Mesalamed-800 TABLETS

Each Delayed Release Tablet Contains: Mesalamine IP 800 mg + Excipients q.s.



#### Indication:

 Mesalamine is used to treat and prevent flare-ups of mild to moderately active ulcerative colitis (an inflammatory bowel disease). It works inside the bowels to reduce inflammation and other symptoms of the disease. This medicine is available only with your doctor's prescription.

#### Dosage:

Directed by the physician

# Mesalamed-1200 TABLETS

Each Delayed Release Tablet Contains: Mesalamine IP 1200 mg + Excipients q.s.



#### Indication:

Mesalamine is used to treat and prevent flare-ups of mild to moderately active ulcerative colitis (an inflammatory bowel disease). It works inside the bowels to reduce inflammation and other symptoms of the disease. This medicine is available only with your doctor's prescription.

#### Dosage:

Directed by the physician

# Methycomed TABLETS

Each film coated tablet contains: Methylcobalamin IP 500 mcg + Excipients q.s.



#### Indication:

 It is prescribed for people suffering from a deficiency of vitamin B12. The aim of this vitamin is to help with the proper functioning of the brain and nerves as well as the production of red blood cells. Methylcobalamin helps treat the deficiency of vitamin B12 by producing a substance known as "myelin".



# Metret 25XL TABLETS

Each film coated extended release tablet contains: Metoprolol Succinate IP 23.75 mg Eq. to Metoprolol Tartrate 25 mg + Excipients q.s.



#### Indication:

- ▶ Yields greater BP reduction and higher response rates vs indivual monotherapies
- ▶ Greater improvement in anginal symptoms vs beta blocker monotherapy

#### Dosage:

Directed by the physician

# Metret 50XL TABLETS

Each film coated extended release tablet contains: Metoprolol Succinate IP 47.5 mg Eq. to Metoprolol Tartrate 50 mg + Excipients q.s.



#### Indication:

- ▶ Yields greater BP reduction and higher response rates vs indivual monotherapies
- Greater improvement in anginal symptoms vs beta blocker monotherapy

#### Dosage:

Directed by the physician

# Metsum-500 TABLETS

Each uncoated tablet contains: Metformin Hydrochloride IP 500 mg + Excipients q.s.



- Decrease platelet aggregation, enhance fibrionlysis
- ▶ Better control in HbAlc by 3.4%

Dosage:

Directed by the physician

# Metsum 500SR TABLETS

Each uncoated sustained release tablet contains: Metformin Hydrochloride IP 500 mg + Excipients q.s.



Decrease platelet aggregation, enhance fibrionlysis

Better control in HbA1c by 3.4%



# Metsum-850 TABLETS

Each Uncoated tablet contains: Metformin Hydrochloride IP 850 mg + Excipients q.s.



#### Indication:

- ► Decrease platelet aggregation, enhance fibrionlysis
- Better control in HbAlc by 3.4%

#### Dosage:

Directed by the physician

# Mira-BG 25 TABLETS

Each film coated extended release tablet contains: Mirabegron 25mg, Excipients q.s.



#### Indication:

- Effectively reduces the severity of overactive bladder and related symptoms.
- ► Convenient once-daily treatment

#### Dosage:

Directed by the physician

# Metsum-1000 SR TABLETS

Each uncoated sustained release tablet contains: Metformin Hydrochloride IP 1000 mg + Excipients q.s.



#### Indication:

- Decrease platelet aggregation, enhance fibrionlysis
- Better control in HbA1c by 3.4%

#### Dosage:

Directed by the physician

# Mira-BG 50 TABLETS

Each film coated extended release tablet contains: Mirabegron 50mg, Excipients q.s.



#### Indication:

- Effectively reduces the severity of overactive bladder and related symptoms.
- ► Convenient once-daily treatment

#### Dosage:



# Methotreat-2.5 TABLETS

Each uncoated tablet contains: Methotrexate Tablets IP 2.5mg.



 Methotrexate is a medication that treats inflammatory arthritis, psoriasis and other inflammatory conditions.

#### Dosage:

Directed by the physician

# Methotreat-7.5 TABLETS

Each uncoated tablet contains: Methotrexate Tablets IP 7.5mg.



► Methotrexate is a medication that treats inflammatory arthritis, psoriasis and other inflammatory conditions.

#### Dosage:

Directed by the physician

# Methotreat-5 TABLETS

Each uncoated tablet contains: Methotrexate Tablets IP 5mg.

# 10X10



 Methotrexate is a medication that treats inflammatory arthritis, psoriasis and other inflammatory conditions.

#### Dosage:

Directed by the physician

# Methotreat-10 TABLETS

Each uncoated tablet contains: Methotrexate Tablets IP 10mg.



 Methotrexate is a medication that treats inflammatory arthritis, psoriasis and other inflammatory conditions.



# Naproxum-250 TABLETS

Each Film Coated Tablet Contains: Naproxen Sodium PH EUR 250 mg + Excipients q.s.



#### Indication:

- Nonsteridal anti-inflammatory drug (NSAID)
- Used to reliev pain from various conditions such as headaches, muscles aches, tendnitis, dental pain, and menstrual cramp

#### Dosage:

Directed by the physician

# Naproxum-D TABLETS

#### Each Film Coated Tablet Contains:

Naproxen Sodium PH EUR 500 mg +Domperidone IP 10mg Excipients q.s.



#### Indication:

- Nonsteridal anti-inflammatory drug (NSAID)
- ► Used to reliev pain from various conditions such as headaches, muscles aches, tendnitis, dental pain, and menstrual cramp

#### Dosage:

Directed by the physician

# Naproxum-500 TABLETS

Each Film Coated Tablet Contains: Naproxen Sodium PH EUR 500 mg + Excipients q.s.



#### Indication:

- Nonsteridal anti-inflammatory drug (NSAID)
- Used to reliev pain from various conditions such as headaches, muscles aches, tendnitis, dental pain, and menstrual cramp

#### Dosage:

Directed by the physician

# Nebihit-2.5 TABLETS

Each Uncoated tablet contains: Nebivolol Hydrochloride IP Eq. to Nebivolol 2.5 mg + Excipients q.s.



#### Indication:


 Nebivolol is used alone or in combination with other medications to treat high blood pressure. Nebivolol is in a class of medications called beta blockers. It works by relaxing blood vessels and slowing heart rate to improve blood flow and decrease blood pressure.

#### Dosage:



#### Nebihit-5 TABLETS

Each Uncoated tablet contains: Nebivolol Hydrochloride IP Eq. to Nebivolol 5 mg + Excipients q.s.



#### Indication:

Nebivolol is used alone or in combination with other medications to treat high blood pressure. Nebivolol is in a class of medications called beta blockers. It works by relaxing blood vessels and slowing heart rate to improve blood flow and decrease blood pressure.

#### Dosage:

Directed by the physician

### Nicosum-10 TABLETS

Each Uncoated Tablet Contains: Nicorandil IP 10 mg + Excipients q.s.



#### Indication:

▶ Nicorandil is a medicine used to treat and reduce chest pain caused by angina. It works by relaxing and widening your blood vessels and increasing the blood and oxygen supply to your heart. Your doctor will usually prescribe nicorandil when other heart medicines have not worked or are not suitable for you.

#### Dosage:

Directed by the physician

# Nicosum-5 TABLETS

Each Uncoated Tablet Contains: Nicorandil IP 5 mg + Excipients q.s.



#### Indication:

▶ Nicorandil is a medicine used to treat and reduce chest pain caused by angina. It works by relaxing and widening your blood vessels and increasing the blood and oxygen supply to your heart. Your doctor will usually prescribe nicorandil when other heart medicines have not worked or are not suitable for you.

#### Dosage:

Directed by the physician

### Nicorole-1 TABLETS

Each Uncoated tablet contains: Nicoumalone IP 1mg Excipients q.s.



#### Indication:

 Acenocoumarol (also called nicoumalone) is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood. **Dosage:** 



# Nicorole-2 TABLETS

Each Uncoated tablet contains: Nicoumalone IP 2mg Excipients q.s.



#### Indication:

▶ Acenocoumarol (also called nicoumalone) is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood. **Dosage:** 

Directed by the physician

# Nicorole-4 TABLETS

Each Uncoated tablet contains: Nicoumalone IP 4mg Excipients q.s.



#### Indication:

 Acenocoumarol (also called nicoumalone) is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood. Dosage:

Directed by the physician

### Nicorole-3 TABLETS

Each Uncoated tablet contains: Nicoumalone IP 3mg Excipients q.s.



#### Indication:

 Acenocoumarol (also called nicoumalone) is used to prevent and treat harmful blood clots, reducing the risk of a stroke or heart attack. This medication helps to keep blood flowing smoothly in your body by decreasing the amount of clotting proteins in the blood. **Dosage:** 

Directed by the physician

# Nitrofuromed-100SR TABLETS

Each Uncoated Tablet Contains: Nitrofurantoin IP 100 mg + Excipients q.s.



#### Indication:

▶ Nitrofurantoin is used to treat urinary tract infections. This medicine is an antibiotic. It works by killing bacteria or preventing their growth. However, this medicine will not work for colds, flu, or other virus infections.

#### Dosage:



 $1 \times 30$ 

M.R.P.₹ 240/-

# Nitro-Gilsum 2.6 TABLETS

Each uncoated controlled release tablet contains: Diluted NItroglycerin IP Eq. to Nitroglycerin 2.6 mg (In a controlled release system) + Excipients q.s.



#### Indication:

► For prophylaxis & long-term Management of Angina pectoris, heart failure and MI

#### Dosage:

Directed by the physician

# Oxypine-300 TABLETS

Each Film coated tablet contains: Ofloxacin IP 200 mg.



#### Indication:

Oxcarbazepine is used alone or together with other medicines in the treatment of epilepsy to control partial seizures. It works in the brain to prevent seizures. However, this medicine will not cure epilepsy and will only control seizures for as long as you continue to take it.

#### Dosage:

Directed by the physician

# Nitro-Gilsum 6.4 TABLETS

Each uncoated controlled release tablet contains: Diluted NItroglycerin IP Eq. to Nitroglycerin 6.4 mg (In a controlled release system) + Excipients q.s.



#### Indication:

 For prophylaxis & long-term Management of Angina pectoris, heart failure and MI

#### Dosage:

Directed by the physician

# Olnipine-5 TABLETS

Each uncoated tablet contains : Olanzapine Tablets IP 5mg Excipients q.s.



#### Indication:

 Olanzapine is used to treat schizophrenia. It may also be used alone or with other medicines (eg, lithium or valproate) to treat mania or mixed episodes that is part of bipolar disorder ( manic-depressive illness).

#### Dosage:



# Olnipine-10TABLETS

Each uncoated tablet contains : Olanzapine Tablets IP 10mg Excipients q.s.



#### Indication:

 Olanzapine is used to treat schizophrenia. It may also be used alone or with other medicines (eg, lithium or valproate) to treat mania or mixed episodes that is part of bipolar disorder ( manic-depressive illness).

#### Dosage:

Directed by the physician

# Phensum-100 TABLETS

#### Phenytoin 100MG TAB

Each film coated tablet contains: Phenytoin Sodium IP 100 mg + Excipients q.s.



#### Indication:

 phenytoin is used to control certain type of seizures, and to treat and prevent seizures that may begin during or after surgery to the brain or nervous system.
 Phenytoin is in a class of medications called anticonvulsants. It works by decreasing abnormal electrical activity in the brain.

#### Dosage:

Directed by the physician

### PG-Win-75 TABLETS

(Pregabalin IP 75 mg + Excipients q.s.

### 10x10



#### Indication:

 Pregabalin is used to treat epilepsy and anxiety. It's also taken to treat nerve pain. Nerve pain can be caused by different conditions including diabetes and shingles, or an injury.

#### Dosage:

Directed by the physician

### Prazowin-XL 2.5 TABLETS

Each uncoated Sustained release tablet contains : prazosin hydrochloride IP 2.5mg Excipients q.s.



#### Indication:

Prazosin is used with or without other medications to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Prazosin belongs to a class of medications called alpha blockers. It works by relaxing and widening blood vessels so blood can flow more easily.

#### Dosage:





# Prazowin-XL 5 TABLETS

Each uncoated Sustained release tablet contains : prazosin hydrochloride IP 5mg Excipients g.s.

# 10x1x10



#### Indication:

Prazosin is used with or without other medications to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Prazosin belongs to a class of medications called alpha blockers. It works by relaxing and widening blood vessels so blood can flow more easily.

#### Dosage:

Directed by the physician

# Ramiril-2.5 TABLETS

Each film coated tablet contains: Ramipril IP 2.5 mg + Excipients q.s.



#### Indication:

- ▶ Significant reduction in risk of total by 24% VS placebo
- ▶ Significant reduce risk of cardicovascular events by 25% placbo
- Significant reduction by 22% in composite o mi, stroke or cv death in high risk patients vs placebo

#### Dosage:

Directed by the physician

# Prasugrate-10 TABLETS

Each film Coated tablet Contains: Prasugrel 10mg Excipients q.s.

#### 20x30



#### ndication:

Prasugrel is used along with aspirin to prevent serious or life-threatening problems with the heart and blood vessels in people who have had a heart attack or severe chest pain and have been treated with angioplasty (procedure to open the blood vessels that supply blood to the heart).

#### Dosage:

Directed by the physician

### Ramiril–5 TABLETS

Each film coated tablet contains: Ramipril IP 5mg + Excipients q.s.



#### Indication:

- ► Significant reduction in risk of total by 24% VS placebo
- ▶ Significant reduce risk of cardicovascular events by 25% placbo
- Significant reduction by 22% in composite o mi, stroke or cv death in high risk patients vs placebo

#### Dosage:



# Ranozine-500 TABLETS

Each Film Coated Tablet Contains: Ranolazne 500mg. (Sustained Release) Excipients q.s.



#### Indication:

- Ranolazine is a piperazine derivative
- Treats chronic angina (chest pain)
- ► Has similar efficacy in younger and elderly patients.

#### Dosage:

Directed by the physician

#### Rosvasum-20 TABLETS

Each film coated tablet contains: Rosuvastatin Calcium IP Eq. to Rosuvastatin 20 mg + Excipients q.s.



#### Indication:

Rosuvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.

Directed by the physician

#### Rosvasum–10 TABLETS

Each film coated tablet contains: Rosuvastatin Calcium IP Eq. to Rosuvastatin 10 mg + Excipients g.s.



- ▶ Rosuvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It belongs to a group of drugs known as "statins." It works by reducing the amount of cholesterol made by the liver.

#### Dosage:

Directed by the physician

#### Repaglide 0.5 TABLETS

Each film coated tablet contains: Repaglinidine 0.5mg+ Excipients q.s.



#### Indication:

Repaglinide is used together with diet and exercise to treat high blood sugar (glucose) levels in patients with type 2 diabetes. Repaglinide works by causing your pancreas to release more I nsulin into the blood stream. This medicine does not help patients who have insulin-dependent or type 1 diabetes.





# Repaglide 1

TABLETS

Each film coated tablet contains: Repaglinidine 1mg+ Excipients q.s.



#### Indication:

▶ Repaglinide is used together with diet and exercise to treat high blood sugar (glucose) levels in patients with type 2 diabetes. Repaglinide works by causing your pancreas to release more I nsulin into the blood stream. This medicine does not help patients who have insulin-dependent or type 1 diabetes.

Directed by the physician

# Seveler-800 TABLETS

Each film coated tablet contains: Sevelamer Carbonate 800mg. Excipients q.s.



- Hyperphosphatemia
- ▶ Hypocalcemia
- Hyperparathyroidism

#### Dosage:

Directed by the physician

# **Repaglide 2**

TABLETS

Each film coated tablet contains: Repaglinidine 2mg + Excipients q.s.



#### Indication:

▶ Repaglinide is used together with diet and exercise to treat high blood sugar (glucose) levels in patients with type 2 diabetes. Repaglinide works by causing your pancreas to release more I nsulin into the blood stream. This medicine does not help patients who have insulin-dependent or type 1 diabetes.

Directed by the physician

# Seveler-400 TABLETS

Each film coated tablet contains: Sevelamer Carbonate 400mg. Excipients q.s.





# Sitasure-50 TABLETS

Each film coated tablet contains:

sitagliptin phosphate monohydrate ip eq. to sitagliptin 50mg



#### Indication:

- ▶ Increase concentration of GLP-1 & GIP therapy increasing & Prolonging the action of these hormones.
- ▶ Increase insulin release & decrease glucagon levels in the circulation.

#### Dosage:

Directed by the physician

#### Sitasure–M 500 TABLETS

Each film coated tablet contains: sitagliptin phosphate monohydrate ip eq. to sitagliptin 50mg, metformin hydrochloride ip 500mg.



#### Indication:

- ▶ Increase concentration of GLP-1 & GIP therapy increasing & Prolonging the action of these hormones.
- ▶ Increase insulin release & decrease glucagon levels in the circulation.

#### Dosage:

Directed by the physician

# Sitasure–100 TABLETS

#### Each film coated tablet contains:

sitagliptin phosphate monohydrate ip eq. to sitagliptin 100mg



#### Indication:

- ► Increase concentration of GLP-1 & GIP therapy increasing & Prolonging the action of these hormones.
- ▶ Increase insulin release & decrease glucagon levels in the circulation.

#### Dosage:

Directed by the physician

### Sitasure–M 1000 TABLETS

Each film coated tablet contains: sitagliptin phosphate monohydrate ip eq. to sitagliptin 50mg , metformin hydrochloride ip 1000mg.



#### Indication:


- ▶ Increase concentration of GLP-1 & GIP therapy increasing & Prolonging the action of these hormones.
- ▶ Increase insulin release & decrease glucagon levels in the circulation.

#### Dosage:



# Sertasum-50 TABLETS

#### Each uncoated tablet contains : Sertraline IP 50mg. Excipients q.s.



certain actions over and over), panic attacks (sudden, unexpected attacks of extreme fear and worry about these attacks), posttraumatic stress disorder (disturbing psychological symptoms

#### Dosage:

Directed by the physician

# Solifenasum–5 TABLETS

Each Film Coated Tablet Contains: Solifenacin Succinate 5 mg + Excipients q.s.



#### Indication:

 Solifenacin is used to treat bladder problems, including neurogenic detrusor overactivity and symptoms of an overactive bladder, such as incontinence (loss of bladder control), a strong need to urinate right away, or a frequent need to urinate.

#### Dosage:

Directed by the physician

# Sodicomed-500 TABLETS

Each Film Coated Tablet Contains: Sodium Bicarbonate IP 500 mg + Excipients q.s.



#### Indication:

Sodium bicarbonate, also known as baking soda, is used to relieve heartburn, sour stomach, or acid indigestion by neutralizing excess stomach acid. When used for this purpose, it is said to belong to the group of medicines called antacids. It may be used to treat the symptoms of stomach or duodenal ulcers.

#### Dosage:

Directed by the physician

# Spirofit-25 TABLETS

Each Uncoated Tablet Contains: Spironolactone IP 25 mg + Excipients q.s.

# 10x10





#### Indication:

Spironolactone is used in combination with other medicines to treat high blood pressure (hypertension) and heart failure. Lowering blood pressure can reduce the risk of strokes and heart attacks. Spironolactone is also used to lessen the need for hospitalization for heart failure.

#### Dosage:



# Spirofit-50 TABLETS

Each Uncoated Tablet Contains: Spironolactone IP 50 mg + Excipients q.s.



#### Indication:

▶ Spironolactone is used in combination with other medicines to treat high blood pressure (hypertension) and heart failure. Lowering blood pressure can reduce the risk of strokes and heart attacks. Spironolactone is also used to lessen the need for hospitalization for heart failure.

#### Dosage:

Directed by the physician

# Terbihit-250 TABLETS

#### Each uncoated tablet contains: Terbinafine Hydrochloride USP Eq. to Terbinafine 250 mg + Excipients q.s.



#### Indication:

▶ Fungicidal against dermatophytes, moulds and certain fungi highly kerainophilic with a bioavailability of 70% to 80% treatment of choice with in the therapy of onychomycosis because of effectiveness well tolerated with low potential for drug interactions

#### Dosage:

Directed by the physician

# Sulfasin-500 TABLETS

Each enteric-coated tablet contains : Sulfasalazine IP 500 mg

# 10x10



#### Indication:

▶ Sulfasalazine is used to treat ulcerative colitis and Crohn's disease, and other types of inflammatory bowel disease. It can also be used to treat rheumatoid arthritis. It belongs to a group of medicines called aminosalicylates.

#### Dosage:

Directed by the physician

# Telmihit-H TABLETS

Each film coated tablet contains: Telmisartan IP 40 mg +Hydrochlorothiazide IP 12.5 mg + Excipients q.s.



#### Indication:

► Telmisartan and hydrochlorothiazide combination is used to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

#### Dosage:



## Telmihit-40 TABLETS

Each film coated tablet contains: Telmisartan IP 40 mg + Excipients q.s.



### Indication:

Telmisartan is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

### Dosage:

Directed by the physician

## Torsemed-10 TABLETS

Each uncoated tablet contains : Torsemide IP 10 mg Excipients q.s.



### Indication:

Torsemide is used to help treat fluid retention (edema) and swelling that is caused by congestive heart failure, liver disease, kidney disease. It belongs to the group of medicines called loop diuretics (water pills). This medicine works by acting on the kidneys to increase the flow of urine.

### **Dosage:** Directed by the physician

## Telmihit-AM TABLETS

Each film coated tablet contains: Telmisartan IP 40 mg + Amlodipine 5mg Excipients q.s.

	25x10
	M.R.P.₹ 90/-
	41 staldeT aniqibolmA & netresimlaT
5	25x10 Tablets
linihi	Telmisartan & Amlodipine Tablets IP
	Telmihit-AM
ì	

### Indication:

Telmisartan is used alone or together with other medicines to treat high blood pressure (hypertension). High blood pressure adds to the workload of the heart and arteries. If it continues for a long time, the heart and arteries may not function properly.

### Dosage:

Directed by the physician

## Torsemed-20 TABLETS

Each uncoated tablet contains : Torsemide IP 20 mg Excipients q.s.



### Indication:

Torsemide is used to help treat fluid retention (edema) and swelling that is caused by congestive heart failure, liver disease, kidney disease. It belongs to the group of medicines called loop diuretics (water pills). This medicine works by acting on the kidneys to increase the flow of urine.

### Dosage:



## Torsemed-40 TABLETS

Each uncoated tablet contains : Torsemide IP 40 mg Excipients q.s.



### Indication:

Torsemide is used to help treat fluid retention (edema) and swelling that is caused by congestive heart failure, liver disease, kidney disease. It belongs to the group of medicines called loop diuretics (water pills). This medicine works by acting on the kidneys to increase the flow of urine.

### Dosage:

Directed by the physician

## Torsemed-S TABLETS

Each uncoated tablet contains : Torsemide IP 100 mg + Spironolactone 10mg Excipients q.s.



### Indication:

► Torsemide is used to help treat fluid retention (edema) and swelling that is caused by congestive heart failure, liver disease, kidney disease. It belongs to the group of medicines called loop diuretics (water pills). This medicine works by acting on the kidneys to increase the flow of urine.

### **Dosage:** Directed by the physician

## **Torsemed-100** TABLETS

Each uncoated tablet contains : Torsemide IP 100 mg Excipients q.s.



### Indication:

Torsemide is used to help treat fluid retention (edema) and swelling that is caused by congestive heart failure, liver disease, kidney disease. It belongs to the group of medicines called loop diuretics (water pills). This medicine works by acting on the kidneys to increase the flow of urine.

### Dosage:

Directed by the physician

### Thyro-null-25 TABLETS

Each uncoated tablet contains Thyhroxine Sodium IP eq. to Anhydrous Thyroxine Sodium 25mg. Excipients q.s



### Indication:

Thyronull is used to treat an underactive thyroid. It replaces or provides more thyroid hormone, which is normally produced by the thyroid gland. Low thyroid hormone levels can occur naturally or when the thyroid gland is injured by radiation/medications or removed by surgery.

### Dosage:

## TABLETS



1X120

M.R.P.₹ 192.19/-

## Thyro-null-50 TABLETS

Each uncoated tablet contains Thyhroxine Sodium IP eq. to Anhydrous Thyroxine Sodium 50mg. Excipients q.s



### Indication:

Thyronull is used to treat an underactive thyroid. It replaces or provides more thyroid hormone, which is normally produced by the thyroid gland. Low thyroid hormone levels can occur naturally or when the thyroid gland is injured by radiation/medications or removed by surgery.

### Dosage:

Directed by the physician

## Thyro-null-100 TABLETS

Each uncoated tablet contains

Thyhroxine Sodium IP eq. to Anhydrous Thyroxine Sodium 100mg. Excipients q.s



### Indication:

Thyronull is used to treat an underactive thyroid. It replaces or provides more thyroid hormone, which is normally produced by the thyroid gland. Low thyroid hormone levels can occur naturally or when the thyroid gland is injured by radiation/medications or removed by surgery.

#### Dosage:

Directed by the physician

## Thyro-null-75 TABLETS

Each uncoated tablet contains Thyhroxine Sodium IP eq. to Anhydrous Thyroxine Sodium 75mg. Excipients q.s



### Indication:

Thyronull is used to treat an underactive thyroid. It replaces or provides more thyroid hormone, which is normally produced by the thyroid gland. Low thyroid hormone levels can occur naturally or when the thyroid gland is injured by radiation/medications or removed by surgery.


### Dosage:

Directed by the physician

### Tridose – 2

TABLETS

(Each Uncoated Tablet Contains: Trihexyphenidyl Hydrochloride IP 2 mg + Excipients q.s.



### Indication:

Trihexyphenidyl is used alone or together with other medicines to treat Parkinson's disease. By improving muscle control and reducing stiffness, this medicine allows more normal movements of the body as the disease symptoms are reduced.

### Dosage:

## TABLETS



## Trimeta-35 MR TABLETS

Each film coated modified release tablet contains : Trimetazidine Dihydrochloride IP 35mg. Excipients q.s.



### Indication:


Trimetazidine is used to prevent Angina (heart-related chest pain). Trimetazidine is an anti-anginal medication. It decreases the oxygen requirement of the heart by shifting its metabolism from fats to glucose. As a result, the heart works more efficiently.

### Dosage:

Directed by the physician

## Tamsulosum-D TABLETS

Each film coated tablet contains Tamsulosin HCl IP 0.4 mg + Dutasteride 0.5 mg



prostate (benign prostatic hyperplasia or BPH) which include difficulty urinating (hesitation, dribbling, weak stream, and incomplete bladder emptying), painful urination, and urinary frequency and urgency.

## Dosage:

Directed by the physician

## Tamsulosum-0.4 TABLETS

Each film coated tablet contains Tamsulosin Hdyrochloride IP 0.4 mg (As Modified Release) + Excipients q.s.



### Indication:

Tamsulosin is used in men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia or BPH) which include difficulty urinating (hesitation, dribbling, weak stream, and incomplete bladder emptying), painful urination, and urinary frequency and urgency.

### Dosage:

Directed by the physician

### Tiklor-90 TABLETS

Each Film Coated Tablet Contains:3 Ticagrelor IP 90 mg + Excipients q.s.



### Indication:

Ticagrelor is an antiplatelet medicine.
 It makes your blood flow through your veins more easily.

**Dosage:** Directed by the physician



## Tofciti-5 TABLETS

Each Film coated tablet contains: Tofacitinib Citrate IP 5mg eq. to Tofacitinib, Excipients q.s




### Indication:

▶ Reduce joint pain, swelling and further joint damage in rheumtoid arthritis

**Dosage:** Directed by the physician

## Voglimed-0.2 TABLETS

Each uncoated tablet contains: Voglibose IP 0.2 mg + Excipients q.s.



### Indication:

 Voglibose is an alpha-glucosidase inhibitor that is used to treat diabetes mellitus by lowering post-prandial blood glucose levels.
 It helps to delay the absorption of sugar in the body and thereby it reduces the macrovascular complication risks.

### Dosage:

Directed by the physician

## Vildamed-50

TABLETS

Each Uncoated Tablet Contains: Vildagliptin 50 mg + Excipients q.s.

## 10x10



### Indication:

► Vildagliptin is indicated in the treatment of type II diabetes mellitus in adults.

### Dosage:

Directed by the physician

## Voglimed-0.3 TABLETS

Each uncoated tablet contains: Voglibose IP 0.3 mg + Excipients q.s.





### Indication:

 Voglibose is an alpha-glucosidase inhibitor that is used to treat diabetes mellitus by lowering post-prandial blood glucose levels. It helps to delay the absorption of sugar in the body and thereby it reduces the macrovascular complication risks.

### Dosage:



## Valscub-50 TABLETS

(Each coated tabletcontains: Sacubitril sodium eq to sacubitril 24mg + valsartan IP 26 mg + Excipients q.s.



### Indication:

Sacubitril and valsartan combination is used together with other medicines to treat chronic heart failure in adults and cut down on t he number of hospital visits for heart problems. It is also used to t reat symptomatic heart failure in children 1 year of age and older.

### Dosage:

Directed by the physician

## Valscub-200 TABLETS

(Each coated tablet contains: Sacubitril sodium eq to sacubitril 97mg + valsartan IP 103 mg + Excipients q.s.



### Sacubitril and valsartan combination is used together with other medicines to treat chronic heart failure in adults and cut down on t he number of hospital visits for heart problems. It is also used to t reat symptomatic heart failure in children 1 year of age and older.

### Dosage:

Directed by the physician

## Valscub-100 TABLETS

(Each coated tablet contains: Sacubitril sodium eq to sacubitril 49mg + valsartan IP 51 mg+ Excipients q.s.

## 10X14



### Indication:

Sacubitril and valsartan combination is used together with other medicines to treat chronic heart failure in adults and cut down on t he number of hospital visits for heart problems. It is also used to t reat symptomatic heart failure in children 1 year of age and older.

### Dosage:

Directed by the physician

## Vertisum-8 TABLETS

Each uncoated tablet contains: Betahistine Hydrochloride IP 8 mg + Excipients q.s.



 Highly effective with good safety in symptomatic treatment of vestibular and co-ordination disturbances.

**Dosage:** Directed by the physician



## Vertisum-16 TABLETS

Each uncoated tablet contains: Betahistine Hydrochloride IP 16 mg + Excipients q.s.



### Indication:

 Highly effective with good safety in symptomatic treatment of vestibular and co-ordination disturbances.

### Dosage:

Directed by the physician

## Zyciryk-100 TABLETS

Each Uncoated Tablet Contains: Allopurinol IP 100 mg + Excipients q.s



### Indication:

Allopurinol is used to prevent or lower high uric acid levels in the blood. It is also used to prevent or lower excess uric acid levels caused by cancer medicines or in patients with kidney stones. A high uric acid level can cause gout or gouty arthritis (joint pain and inflammation).

### Dosage:

Directed by the physician



10x10

M.R.P.₹ 21.17/-

10x10

## CAPSULES



## Dabitran-110 CAPSULES

Each Hard Gelatin Capsule Contains: Dabigatran Etexilate Mesylate 126.83mg Eq. to Dabigatran Etexilate 110 mg+ Excipients q.s.



### Indication:

 Dabigatran is used to prevent stroke and harmful blood clots (such as in your legs or lungs) if you have a certain type of irregular heartbeat (atrial fibrillation).

### Dosage:

Directed by the physician

## Omeesum-D CAPSULES

Each hard gelatin capsule contains: Omeprazole IP 20mg (As enteric- coated pellets) + Domperidone IP 10 mg (As pellets) + Excipients q.s.



► In Gastro oesophageal reflux diseas

▶ In gastro ulcers

Dosage:

Directed by the physician

### Omeesum

Each hard gelatin capsule contain : Omeprazole I.P 20MG

## **20x15**



### Indication:

In Gastro oesophageal reflux diseas
In gastro ulcers

### Dosage:

Directed by the physician

## Rosvasum-AP 10 CAPSULES

Each Hard Gelatin Capsule Contains:Rosuvastatin Calcium IP Equivalent to Rosuvastatin 10 mg (As Granules) + Aspirin IP 75 mg (As gastro-resistanttablet) + Excipients q.s.



### Indication:

Rosuvastatin+Aspirin is a combination of blood-thinning agent and cholesterol-lowering agent, primarily used to prevent myocardial infarction (heart attack), stroke or peripheral vascular disease. It is also used to lower abnormally elevated cholesterol or fat levels (hyperlipidaemia or dyslipidaemia) in the body.

### Dosage:

## CAPSULES



## Rosvasum-AP 20 CAPSULES

Each Hard Gelatin Capsule Contains:Rosuvastatin Calcium IP Equivalent to Rosuvastatin 20 mg (As Granules) + Aspirin IP 75 mg (As gastro-resistanttablet) + Excipients q.s.

10x10



selling of the selling selling

Rosuvastatin & Aspirin Capsules

### Indication:

Rosuvastatin+Aspirin is a combination of blood-thinning agent and cholesterol-lowering agent, primarily used to prevent myocardial infarction (heart attack), stroke or peripheral vascular disease. It is also used to lower abnormally elevated cholesterol or fat levels (hyperlipidaemia or dyslipidaemia) in the body.

10x10 Cansules

Dosage:

Directed by the physician

## Rosvasum-Gold 20 CAPSULES

Each Hard Gelatin Capsule Contains:Rosuvastatin Calcium IP Eq.to Rosuvastatin 20 mg (As Granules) +Clopidogrel Bisulphate IP Eq.to Clopidogrel 75 mg + (As Granules) +Aspirin IP 75 mg (As Gastro-resistant tablets IP) + Excipients q.s.)



### Indication:

 Rosuvastatin+clopidogrel+aspirin belongs to a group of antihyperlipidemic and antiplatelets used to treat heart attack, myocardial infarction (blockage of the blood to muscles of the heart), angina (chest pain), and stroke.

#### Dosage:

Directed by the physician

## Rosvasum-Gold 10 CAPSULES

Each Hard Gelatin Capsule Contains:Rosuvastatin Calcium IP Eq.to Rosuvastatin 10 mg (Ås Granules) +Clopidogrel Bisulphate IP Eq.to Clopidogrel 75 mg + (Ås Granules) +Åspirin IP 75 mg (Ås Gastro-resistant tablets IP) + Excipients q.s.)

### 10x10



Rosuvastatin, Aspirin & Clopidogril Capsule Rosuvastatin, Aspirin & Clopidogril Capsule

alle and a clopidoda anges , alle

### Indication:

Rosuvastatin+clopidogrel+aspirin belongs to a group of antihyperlipidemic and antiplatelets used to treat heart attack, myocardial infarction (blockage of the blood to muscles of the heart), angina (chest pain), and stroke.

#### Dosage:

Directed by the physician

## Tacromed-0.5 CAPSULES

Each Hard Geletin Capsule Contains: Tacrolimus Ip 0.5 Mg + Excipients. Q.s



### Indication:

Tacrolimus is used to stop your body from attacking or rejecting your transplanted organ (heart, kidney, liver, or lung). This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

### Dosage:





## Tacromed-0.5 CAPSULES



### Indication:

 Tacrolimus is used to stop your body from attacking or rejecting your transplanted organ (heart, kidney, liver, or lung).
 This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

### Dosage:

Directed by the physician

Alfacalcidol IP 0.25 mcg

## ALPHA-CAL-C CAPSULES

Each Soft Gelatin Capsule Contains



### Indication:

- ▶ Strengthens bones and reduce total fracture rate
- Prevents & treats osteoporosis & osteoarthritis.
- ▶ Increase absorption of calcium & treats Vitamin D deficiency

### Dosage:

Directed by the physician





10x10



## Brohexum SYRUP



## cough s

Directed by the physician

## Lactosum SYRUP

Each 15 ml contains: Lactulose Concerntrate USP Eq. to Lactulose (66%w/v) 10gm + Aqueous base q.s.



### Indication:

- Draw water in into the bowel through osmotic action
- Active water retaining action softens stools

**Dosage:** Directed by the physician

## Cetrizine SYRUP

Each 5ml. Contains: Cetirizine Hydrochloride I.P. 5mg. (colour: sunset yellow FCF in a flavoured syrupy base.






## Indication:

 Cetirizine prevents and treats allergy symptoms, such as red, itchy eyes, sneezing, a runny or stuffy nose, or hives. + It works by blocking histamine, a substance released by the body during an allergic reaction.

### Dosage:





30g.

## Vovemed

OINTMENT

Diclofenac Diethylamine IP 1.16% w/w + (Eq. to Diclofenac Sodium I.P. 1.00% w/w) + + Benzyl Alcohol IP 1.00% w/w (As Preservative) + In Gel Base



### Indication:

Sprains


Strains

## Dosage:

Apply on affected area 3-4 times a day

## Tacromed ointment

(Each gram of ointment contains: Tacrolimus IP 0.10% + Butylated Hydroxy Toluene 0.01% w/w Ointment base q.s.)



Low back pain

Arthritic pain

### Indication:

Tacrolimus topical is used on the skin to treat moderate to severe atopic dermatitis in patients who have received other medicines that have not worked well. Atopic dermatitis is a skin condition where there is itching, redness, and inflammation, much like an allergic reaction.

#### Dosage:

Directed by the physician

## Terbihit

Terbihit Cream (terbinfine Hydrochloride B.p 1 % W/w



### Indication:

Terbinafine is applied to the skin to treat: ringworm of the body (tinea corporis); ringworm of the foot (interdigital and plantar tinea pedis; athlete's foot); ringworm of the groin (tinea cruris; jock itch);

### Dosage:

Apply on affected area 3-4 times a day



## Bimtomet Eye/Ear Drop

Bimatoprost 0.03%w/v +Benzalkonium Chloride Solution I.P. 0.01%v/v (as preservative)



### Indication:

Bimatoprost ophthalmic is used to treat glaucoma (a condition in which increased pressure in the eye can lead to gradual loss of vision) and ocular hypertension (a condition which causes increased pressure in the eye). Bimatoprost is in a class of medications called prostaglandin analogs.

### Dosage:

Directed by the physician

#### Flurifen Eye/Ear Drop

Flubiprofen Opthalmic Solution



### Indication:

► Flurbiprofen eye drops are used to keep the pupils of the eye from getting smaller (miosis) during an eye surgery. This medicine is a topical nonsteroidal anti-inflammatory drug (NSAID). This medicine is available only with your doctor's prescription.

### Dosage:

Directed by the physician

## Dorzofix Eye/Ear Drop

Dorzolamide Hydrochloride IP 2%w/v + Benzalkonium Chloride IP 0.02%v/v



### Indication:

▶ Dorzolamide is used to treat high pressure inside the eye due to glaucoma (open angle-type) or other eye diseases (such as ocular hypertension). Lowering high pressure inside the eye helps to prevent blindness. This medication works by decreasing the amount of fluid within the eye.

Directed by the physician

## Relax Eye/Ear Drop

Hydroxypropyl Methylcellulose IP 0.3% w/v Phenylmecuric nitrate IP 0.01% w/v (As preservative) Sterile Aqueous base q.s.



### Indication:

▶ It is used to relieve dryness and irritation caused by reduced tear flow. It helps prevent damage to the eye in certain eye diseases. Hydroxypropyl methylcellulose may also be used to moisten hard contact lenses and artificial eyes. In addition, it may be used in certain eye examinations.



## Hylosum Eye/Ear Drop

Sodium Hyaluronate B.P. 0.1% w/v + Sodium Perborate B.P. 0.028% w/v (as Preservative) Sterile Aqueous base q.s.



 Increase the stablity of pre-corneal tear film thereby protects the eye surface envionmental aggrvessions

### Dosage:

Directed by the physician

## Latnomet Eye/Ear Drop

Latanoprost IP 0.005% w/v + Benzalkonium Chloride Solution IP 0.02% v/v (as preservative) + Sterile Aqueous Base q.s.



### Indication:

Latanoprost ophthalmic is used to treat glaucoma (a condition in which increased pressure in the eye can lead to gradual loss of vision) and ocular hypertension (a condition which causes increased pressure in the eye). Latanoprost is in a class of medications called prostaglandin analogs.

### Dosage:

Directed by the physician



Ketorolac eye drops is used to treat itching caused by seasonal allergic conjunctivitis (an allergy that occurs at only certain times of the year). It is also used to treat pain, burning, and inflammation of the eye following cataract surgery or corneal refractive surgery.

### Dosage:

Directed by the physician

## Latnomet-T Eye/Ear Drop

Latanoprost IP 0.005% w/v + Benzalkonium Chloride Solution IP 0.02% v/v (as preservative) + Sterile Aqueous Base q.s.



### Indication:

Latanoprost ophthalmic is used to treat glaucoma (a condition in which increased pressure in the eye can lead to gradual loss of vision) and ocular hypertension (a condition which causes increased pressure in the eye). Latanoprost is in a class of medications called prostaglandin analogs.

### Dosage:

## EYE /EAR DROP



## Moxiflosum-D Eye/Ear Drop

Moxifloxacin Hydrochloride IP 0.5%w/v + Dexamethasone Sodium Phosphate 0.1%w/v + Benzalkonium chloride IP 0.02% v/v Sterile aqueous base



## Nepamet Eye/Ear Drop

Nepafenac 0.1%w/v, Sterile Aqueous base q.s.




### Dosage:

Directed by the physician

## Moxiflosum Eye/Ear Drop

Moxifloxacin Hydrochloride IP 0.5%w/v



## Olepep Eye/Ear Drop

Olopatadine Hydrochloride IP eq. to Olopatadine 0.1% w/v + Benzalkonium Chloride Solution (As Preservative) IP 0.02% v/v, Sterile Aqueous base q.s.



► High rates of efficacy in treating ocular itching **Dosage:** 



## Polymet Eye/Ear Drop

Polyethylene Glycol 400 0.4%w/v + Propylene Glycol IP 0.3% w/v + Sterile Aqueous Base q.s.



### Indication:

▶ This medication is used to relieve dry, irritated eyes. Common causes for dry eyes include wind, sun, heating/air conditioning, computer use/reading, and certain medications.

### Dosage:

Directed by the physician

## Tearsum Fresh DS Eye/Ear Drop

(Sodium Carboxymethly- Cellulose I.P 1% w/v Stabilized Oxychloro complex 0.005% w/v



### Indication:

▶ For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun.

▶ May be used as a protectant against further irritation.

### Dosage:

Directed by the physician

## Tearsum Fresh Eye/Ear Drop

Sodium Carboxymethly- Cellulose I.P 0.5% w/v Stabilized Oxychloro complex 0.005% w/v



- ▶ For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun.
- May be used as a protectant against further irritation.

### Dosage:

Directed by the physician

## Travomet Eye/Ear Drop

Travoprost IP 0.004% w/v + Benzalkonium Chloride Solution IP 0.02% v/v (as preservative) + Sterile Aqueous Base q.s.



Dosage: Directed by the physician



## Travomet-T Eye/Ear Drop

Travoprost IP 0.004%w/v Timolol Maleate IP eq. to Timolol 0.5% w/v Benzalkonium Chloride Solution IP 0.02% v/v (as preservative) Sterile Aqueous base q.s.



This combination product is used to treat high pressure inside the eye due to glaucoma (open-angle type) or other eye diseases (such as ocular hypertension). Lowering high pressure inside the eye helps to prevent blindness.

### Dosage:

Directed by the physician

## Budsosum-0.5 Resputes



FOR INHALATION ONLY PRESERVATIVE FREE

A



### Indication:

Budesonide suspension (liquid) for oral inhalation (Pulmicort Respules) is used in children 12 months to 8 years of age. Budesonide belongs to a class of medications called corticosteroids. It works by decreasing swelling and irritation in the airways to allow for easier breathing.

#### Dosage:

Directed by the physician

## Timolol Eye/Ear Drop

Timolol Maleate I.P. eq. to timlol 0.5% w/v, + Benzalkonium Chloride Solution I.P. 0.02% v/v (as Preservative), Sterile Aqueous base q.s.



### **Dosage:** Directed by the physician

### Dousu-Lin Resputes

Each 2.5ml Contains: Levosalbutamol 1.25mg Ipratropium Bromide IP Eq. To Ipratropium Bromide (anhydrous) 500mcg In An Isotonic Solution Q.s.ychloro Complex 0.005% W/v



### Dosage:





Pipramed-TZ Injection

Each Vial Contains: Piperacillin Sodium IP (Sterile) eq. to Piperacillin


4g + Tazobactum Sodium IP (Sterile) eq. to Tazobactam 0.5g

## Fursocort Respules





Each 1 ml injection contains: diclofanac Sodium 75 mg in an aquas base





M.R.P.₹ 28.30/-

### Indication:

Diclofenac injection is a nonsteroidal anti-inflammatory drug (NSAID) used to treat adults with mild to moderate pain. It is used alone or given with other opioid (narcotic) pain medicines to treat adults with moderate to severe pain. This medicine is to be given only by or under the direct supervision of your doctor.

### Dosage:

# **Upcoming Products**

Citicoline tab 1 Citicoline piracetam tab 2 Povidine iodine ointment 3 Prucalopride 2 mg 4 Perindopril 4 mg 5 Perindopril 8 mg 6 Budosunide 1 mg respule 7 **Clotrimazole** ointment 8 Amitriptyline 10mg 9 Amitriptyline 25 mg 10 Nortryptiline 10 mg 11 Nortryptiline 25 mg 12 Saxagliptin 5 mg 13 Saxagliptin 2.5 mg + metformin 500 mg 14 Saxagliptin 2.5 mg+ metformin 1000 mg 15 Febofibrate 145 16 Febofibrate 160 17 Febofibrate 200 18 Moxinidine 0.2 mg Moxinidine 0.3 mg 19



## MEDISUM BIOTECH PVT. LTD.

Regd office : E-5/16, Krishna nagar, Delhi 110051, BDO - 57, Vastushree complex market yard, Pune - 411037, Maharashtra Contact : 011-45502240 web : www.medisumbiotech.com | E-mail : medisumbiotech@gmail.com